

Annual Methodological Archive Research Review

<http://amresearchreview.com/index.php/Journal/about>

Volume 3, Issue 5 (2025)

INCIDENCE AND MORPHOLOGICAL CHARACTERISTICS OF PLANT PARASITIC NEMATODES IN CITRUS ORCHARDS

Madieha Ambreen¹, Fawad Khan², Samina Yasmin³, Farkhanda Manzoor⁴

Article Details

Keywords:

¹Madieha Ambreen

Assistant Professor, School of Biology (Botany Section), Minhaj University Lahore, Pakistan

²Fawad Khan

Medical Entomologist, Entomology Department, Faculty of Chemical and Life Sciences, Abdul Wali Khan University Mardan, Khyber Pakhtunkhwa, Pakistan

³Samina Yasmin

Lecturer, Department of Zoology, Hazara University, Mansehra, Khyber Pakhtunkhwa, Pakistan

⁴Farkhanda Manzoor

Dean of Scientific Research and Development, Minhaj University, Lahore, Pakistan

ABSTRACT

This study investigates the incidence and population densities of plant parasitic nematodes in citrus orchards across three localities: Palai Katlang, Rabbot Dir L, and Dargai Jabbon. The highest disease incidence (70%) was observed in Palai Katlang, followed by Rabbot Dir L (50%) and Dargai Jabbon (40%). Palai Katlang also exhibited the highest nematode population density, particularly in samples S8 (90%) and S9 (85%). Soil analyses revealed that all three locations had similar silt-loam textures, with Palai Katlang having the highest pH (8.94) and electrical conductivity (228 μ S). The presence of *Tylenchulus semipenetrans* and *Radopholus similis*, the main culprits of citrus nematode diseases, was confirmed, with severe root stunting and dieback observed in Palai Katlang. The study underscores the role of soil conditions, such as pH and texture, in influencing nematode survival and density. Recommendations for controlling nematode populations emphasize Integrated Disease Management (IDM) practices, including regular soil sampling and exploration of resistant citrus varieties. Further research is needed to develop effective control strategies to minimize the impact of nematodes on citrus crops.

Introduction

Citrus is one of the most common genera of flowering plants and belongs to (the family *Rutaceae*). The various species of Citrus are all believed to be native to the subtropical and tropical regions of Asia and the Malay Archipelago and to have spread from there to other sections of the world. Citrus has been cultivated through the ages and in some pretty remote places (*Perez & Marvin, 2015*). Citrus is a rich source of sugar, vitamin C, minerals like magnesium, calcium, and organic acid (*Duncan & Cohn, 2005*). In terms of international trade, Citrus is the highest-value fruit. It is grown in more than 125 countries in a belt within 35°latitude south or north of the equator. Around 70% of the world's total citrus production is grown in the Northern Hemisphere and the United States. Brazil is also one of the biggest citrus-producing countries. Citrus supports the world's biggest agriculture industries (*Duncan & Cohn, 2005*). Pakistan is the sixth biggest producer of oranges and kinnow mandarin, and one of the chief citrus-producing countries in the world (*MINFAL, 2005*). Dubai, Saudi Arabia, Kuwait, Qatar, Oman, the Netherlands, Indonesia, Singapore, Malaysia, and the UK are the major Pakistani markets of Kinnow. About ninety-five percent of the Citrus known in the world is provided by Pakistan. In Pakistan, Citrus is produced on an area of 193,985 hectares with an annual production of 2,001,685 tonnes. In Khyber Pakhtunkhwa, citrus fruits were grown on an area of 2,970 hectares with a total production of 24,355 tonnes. Similarly, the land under cultivation of Citrus in Sindh, Balochistan, and Punjab is 5,140, 1,395, and 183,296 hectares at the production rate of 31,394, 6,920, and 1,930,082 tonnes, respectively (*Anonymous, 2013*). A wide range of plant-parasitic nematodes has been associated with the citrus rhizosphere, but only a few species cause damage to trees (*Verdejo-Lucas & McKenry, 2004*). These include migratory endoparasites (lesion and burrowing nematodes), sedentary endoparasites (citrus root-knot nematodes), and several other ectoparasites that cause serious damage to Citrus. The most common nematodes affecting citrus trees are citrus nematodes and *Tylenchus semipenetrans* (*Cobb, 1913*). The nematode causes a "Slow Decline" of Citrus all around the world and limits citrus fruit production under a wide range of environmental conditions (*Duncan, 2005*). All varieties of Citrus are attacked, but rootstocks like trifoliate orange (*Poncirus trifoliata*) are highly resistant to citrus nematode attack. Others, such as Troyer and Carrizo citrange, are moderately tolerant, and some, like sweet oranges, are highly susceptible (*Halbert, 2012*). Citrus nematodes are ectoparasitic and feed on root tissues from outside the plant. With their hindquarters buried in the soil, the larvae pierce the roots to feed, usually just behind the growing tips. Feeding on the roots does not kill the citrus tree. However, the root's capacity to carry water and nutrients is impeded, and yields are noticeably reduced early in the life of the tree. Nematode damage of the roots also promotes the entry of secondary diseases (*Godwin et al., 2007*). Citrus nematodes are most active in the warmer months, from September to April. It is not easy to diagnose these hidden nematodes of Citrus. Spreading decline is another important disease of Citrus caused by *Radopholus similis* (*Cobb, 1893*) *Thorne, 1949*. This nematode caused severe losses in Citrus in central Florida (*Duncan, 2005*). Symptoms in the Citrus included severe necrosis and root destruction (*Machon & Bridge, 1996*). Other species of nematodes, i.e., *Pratylenchus coffeae* (*Zimmermann, 1898*), *Xiphenema* spp., *Paratrichodorus* spp. (*Topham PB. 1985*), *Belonolaimus longicaudatus* (*Mulrooney RP, 2010*) and *Helicotylenchus* spp. (*Whittaker LM. 1984*) have been

reported in citrus orchards. In Pakistan, information on plant-parasitic nematodes attacking citrus orchards is very limited. No reliable data on incidence, occurrence, frequency distribution, and losses incurred by parasitic nematodes infesting citrus orchards are available in the Khyber Pakhtunkhwa province. The present study was undertaken to estimate the incidence and to identify major parasitic nematodes of Citrus in selected localities. Keeping in view the importance of plant parasitic nematodes, a survey was conducted in citrus-growing orchards at Dargai Jabbon, Palai Katlang, and Rabbot Dir Lower to achieve the following objectives.

MATERIALS AND METHODS

A comprehensive survey of citrus orchards in selected localities was carried out. Three sites, i.e., Palai Babozai, Dargai Jabbon, and Rabbot Dir Lower, were selected for collecting roots and soil samples. Samples were collected from citrus orchards following an atriangular design (Zarina *et al.*, 2015). From every site, ten trees were selected. Citrus orchards from selected localities. (Palai katlang) Moreover, Dargai Jabbon Farm was surveyed. Soil sample (each 10 g) was collected with the help of a soil auger up to a depth of approximately 15-20cm (Southey, 1986). Root samples (each 50 g) were also collected from the same citrus orchards from which soil cores were collected. Ten soil and root samples were collected from each orchard. Samples were randomly taken with the help of a trowel and auger. Debris was removed from the upper 15 cm of soil, and samples were taken from 25-30 cm soil depth in the rhizosphere of the diseased and healthy trees at a distance of 60 to 90 cm away from the tree trunk. A composite of 100g of soil with roots was taken in polythene bags, properly labeled, and brought to the laboratory of Plant Pathology for further study. Juveniles and adults were extracted from the soil by the Whitehead and Hemming method (Whitehead & Hemming, 1965). Fifty grams (50 g) of powder soil was placed over tissue paper in a sieve. The sieve was placed in a steel tray. About two hundred milliliters of water were added to the tray from the corner was added until the soil was properly moistened. The sample was left for 24 hours at room temperature. Nematodes extracted from soil samples were counted under a stereomicroscope at 60X magnification, and their population density of nematodes was recorded (Southey, 1986). Citrus roots were washed carefully with tap water to remove soil particles and then air dried. Feeder roots were soaked in a beaker having 180- 200 mL of distilled water for about 24-48 hours. Roots were washed and cut into 1.0 cm pieces. Fifty grams (50 g) of roots were placed in Petri plates containing distilled water. Plates were incubated for 24 hours at room temperature, and nematodes that crawled out of roots were observed under a Stereomicroscope, and their density was calculated at 60X magnification (Southey, 1986). Major nematodes extracted from citrus roots and rhizosphere were identified at 40X magnification using the key (Cohn, E, 1969). Male and female nematodes were identified based on their morphological characteristics. Parasitic nematode genera associated with citrus roots were identified, and morphological features were observed and compared with the key of (Albert *et al.*, 2010). The percentage of sand, silt, and clay in the inorganic fraction of soil was measured by the Hydrometer Method (Bouyoucos, 1962). Fifty (50g) of soil, clear of debris and gravel, was taken in a special dispersion cup, and 10 ml of dispersing solution was added to it. The cup was then filled with water and placed on a stirrer for 5 minutes. The treated soil was transferred

to a 1000 ml glass cylinder and filled with distilled water up to the 1000 ml mark. A hydrometer and thermometer were placed in it. After 40 seconds, the first hydrometer value and temperature were recorded. Then, the suspension was re-shaken and left over for 2 hours. After 2 hours, the ^{second} hydrometer value and temperature were recorded. 0.3 was added to the readings of the samples for every 1 °C above 20 °C and subtracted 0.3 for every 1 °C below 20 °F. (Bouyoucos, 1962). The Silt, Sand, and Clay percentage was calculated as follows;

40 sec reading

Silt + Clay = _____ X 100

weight of sample

Corrected value at 2 hrs

Clay = _____ X 100

weight of sample

Sand = 100- Silt + Clay

Measurements of electrical conductivity (EC) and pH were determined on a saturation extract of soil or supernatant liquid of a 1:2 soil water suspension. A 15 g soil sample was shaken with 30 mL of distilled water in 250 mL conical flasks for 10 minutes. After that, the suspension was passed through filter paper to remove the soil particles. The EC was then measured by the electrical conductivity meter and pH by the pH meter (Hendershot *et al.*, 1993).

RESULTS

The incidence of citrus nematodes across three localities, namely Palai Katlang, Rabbot Dir L, and Dargai Jabbon, was observed in various percentages, with Palai Katlang showing the highest disease incidence at 70%, followed by Rabbot Dir L at 50%, and Dargai Jabbon at 40%. The population density of citrus nematodes, measured in different samples, was highest in Palai Katlang, with S8 showing a 90% density, followed by S9 at 85% and S6 at 80%. In contrast, the nematode density in Rabbot Dir L was highest in sample S5, with a 55% population density. In Dargai Jabbon, the highest density was recorded in sample S9 at 39%, followed by S10 (33%) and S1 (30%). When combining the mean population densities, Palai Katlang recorded the highest mean population density of 599, followed by Rabbot Dir L at 460 and Dargai Jabbon at 243. In terms of nematode density in citrus roots, the highest densities were recorded in S1 of Palai Katlang and Dargai Jabbon. The soil properties, including pH, electrical conductivity (EC), and texture, were also analyzed. Palai Katlang had a soil pH of 8.94 with an EC of 228 µS and a silt-loam texture. Rabbot Dir L had the highest soil pH at 8.98, with an EC of 220 µS and a similar silt-loam texture. Dargai Jabbon exhibited the lowest soil pH at 8.77 with an EC of 206 µS and also had a silt-loam texture. The soil samples from all three locations showed a similar silt-loam texture with varying percentages of silt, sand, and clay, contributing to the overall soil structure. These findings indicate that the highest nematode population densities were observed in silt-loam textured soils, especially in Palai Katlang, which also exhibited the highest disease incidence.





Table pH, electrical conductivity (EC), and soil texture values of soil samples collected from citrus orchards at three different selected localities.

.NO	Locality	pH (µ S)	EC	Percent Silt	Percent Sand	Percent Clay	Soil Texture
1	Palai Katlang	8.94	207	62.2	42.0	4.0	Silt loam
2	Rabbot dir L	8.98	228	68.0	33.7	4.2	Silt loam
3	Dargai Jabbon	8.77	206	58.0	38.0	3.4	Silt loam




Symptoms and morphology of plant parasitic nematodes in Citrus

Two major genera of plant-parasitic nematodes were isolated from the citrus rhizosphere and roots. Infected citrus trees showed typical decline symptoms caused by citrus nematodes, *Tylenchulus semipenetrans* (Figure 4.2.1). Infested

roots showed severe stunting and decay. Severely infected trees showed yellowing and thinned canopies, exposure of bare crown limbs, and reduced leaf and fruit size. *Tylenchulus semipenetrans* exhibited sexual dimorphism. Males and females showed differences in their morphology. Figure 4.2.2 shows a typical male nematode of *Tylenchulus semipenetrans* isolated from citrus roots. The nematode showed a translucent, white-slender vermiform shape under 40X magnification (Fig. 4 .2.2a). The nematode possessed a delicate style with small knobs.

A single elongated testis and dark-paired spicules were observed in adult males. Figure 4.2.2 (b) shows the structure of a typical sedentary female of *T. semipenetrans*. The posterior portion of a mature female was swollen with an elongated neck, whereas its

The head was embedded into root tissues, as shown in the figure. Burrowing nematode *Radopholus similis* exhibits marked sexual dimorphism. Male nematodes possess a raised lip region and a reduced feeding apparatus (stylet and esophagus) because they are not infective. The tail of the male burrowing nematode has a distinctive bursa extending at least two-thirds of the tail length that it uses to clasp the female body during mating (Fig. 4 .2.3.2.3a). Females do not have a raised lip region but do have a heavily sclerotized and thickened framework. The female style is robust, with three distinct knobs. The vulva, the opening of the reproductive system, is located slightly below mid mid-body (Fig. 4.2.3 b). Infected roots of Citrus revealed another genus of plant parasitic nematode, i.e., spiral nematode, *Helicotylenchus* spp. The nematode showed a typical spiral structure, as shown in the figure (4.2.4). Figure 4.2.6 shows the female spiral nematode. The nematode is curved into a spiral shape when relaxed or dead. The vulva of the female is located around 2/3 of the nematode's body length from the anterior terminus, as shown in Figure (4.2.6). The tail of the nematode is asymmetrical. The male is straight, as shown in the figure (). The nematode produced dark necrotic lesions on roots and caused root dieback.



2

Fig. Adult male (a) and female (b) of *Tylenchulus semipenetrans* isolated from citrus roots.



Fig 4.2.3 *Radopholus similis* male head (A) and tail (B), Left. The female of *R. similis* (right) with the head region (A) and full body with vulva [v] near midbody (B). (Internet Picture: Photograph by [Nicholas Sekora](#))




Fig Adult female of *Helicotylenchus* spp (A) isolated from Citrus. The body of the nematode is curved into a spiral shape when the nematode is dead or relaxed. B) Male spiral nematode (*Helicotylenchus* spp.)

DISCUSSION

Plant-parasitic nematodes are widely distributed pests and are abundant in agricultural soils. In the present study, we detected and identified two major genera of plant parasitic nematodes, i.e., *Helicotylenchus* spp. (spiral nematode) and *Tylenchulus semipenetrans*. These plant parasitic nematodes were found associated with Citrus in three different localities, i.e., Dargai Jambon, Palai Katlang, and Rabbot Dir Lower). We found the highest incidence of citrus nematodes in Palai Katlang (district.) and the minimum in the Dargai Jabbon soils. The population densities of citrus nematodes were the greatest in the Palai Katlang locality. Higher nematode density in roots produced severe stunting and dieback symptoms on roots, particularly in root samples that were collected from Palai Katlang. Higher disease incidence and densities of nematodes in the play Katlang locality could be considered due to favorable soil conditions, temperature, and cultivation of susceptible citrus rootstocks. Numerous citrus varieties are cultivated in Pakistan. These include rough lemon, sour oranges, kinnow, sweet oranges, mandarins, and wild oranges. The nematode population depends on the type of host infested. *T. semipenetrans* cause slow declines in all citrus nurseries and orchards but its incidence varies in all citrus varieties.

We reported a higher incidence of citrus nematodes in sweet oranges. Similar results were reported by other researchers who observed a maximum incidence of citrus nematodes in the Feutrells variety, lemon and orange, and a minimum incidence in grapefruit (Khanzada *et al.*, 2007). Some researchers documented that rough lemon and sour oranges contain a maximum number of female citrus nematodes as compared to neutrals and sweet oranges (Ahmad *et al.*, 2007). Soil texture also affects nematode population densities, reproduction factor and movement of nematodes. As a generalization, sandy soils are nematode-loving soils. *T. semipenetrans* and *Helicotylenchus* spp. Do not persist well in dry soil, and population densities on drought-stressed trees decline rapidly (Tsai & Van Gundy, 1988). However, infection and population growth are rapid in localized areas of drought in the rhizosphere if other portions of the root system have adequate water. However, nematodes exist in soils of all textures, some even being favored by a finer texture. Soil pH is a useful measurement because it is a predictor of various chemical activities within the soil. As such, it is also a useful tool in making management decisions in controlling different soil-borne plant pathogens, including plant parasitic nematodes. Soil pH appears crucial to nematode survival. Acid or alkaline soils tend not to favor nematodes. In conclusion, control of plantparasitic nematodes typically requires an integrated approach. However, there are often no or very few chemical options. In field crops, due to the cost of nematicides, their use is not economically justified. Therefore, cultural tactics should often be the first line of defense for the management of plant-parasitic nematodes. The importance of maintaining proper fertility and optimizing soil pH has been emphasized here to reduce the impact of nematodes.

Conclusions:

Nematodes are widely distributed in citrus-growing areas. Three types of nematode genera, viz., citrus nematode, *Tylenchulus semipenetrans*, and *radiophones similis Helicotylenchus* spp (spiral nematode), were detected and identified. The highest percentage of disease incidence (70%) was recorded in Palai katlang, and the minimum disease incidence (40%) was recorded in Dargai jabbon. The highest Population density of nematodes occurred in Palai Katlang citrus orchards.

REFERENCES

- Brown, D. J. F., P. B. Topham. (1985). Morphometric variability between populations of *Xiphinema diversicaudatum* (Nematoda: Dorylaimoidea). *Revue de Nematologie*. 8: 15-26
- Chabrier C., P. Quénéhervé. 2003. Control of the burrowing nematode *Radopholus similis* Cobb on banana: Impact of the banana field destruction method on the efficiency of the following fallow. *Crop Protection*. 22: 121-127
- Chabrier, C., P. Tixier., P. F. Duyck., C. Carles., P. Quénéhervé. 2010. Factors influencing the survivorship of the burrowing nematode, *Radopholus similis* (Cobb.) Thorne in two types of soil from banana plantations in Martinique. *Applied Soil Ecology*. 44: 116-123.
- Cobb, N. A. (1915). *Tylenchulus similis* is the cause of a root disease of Sugar cane and banana. *Journal of Agricultural Research*. 4(6): 561–568.
- Cohn, E. (1969). The occurrence and distribution of species of *Xiphinema* and *Longidorus* in Israel. *Nematologica*. 15:179–192
- Duncan, L. W., D. M. Eissenstat., 1993. Responses of *Tylenchulus semipenetrans* to citrus fruit removal: implications for carbohydrate competition. *Journal of Nematology*. 25(1):7–14.
- Duncan, L. W., M. M., E.-M., (1996). Population changes of *Tylenchulus semipenetrans* under localized versus uniform drought in the citrus root zone. *Journal of Nematology*. 28(3): 360–368.
- Duncan, L.W., P. M. Ashela., J. Ferguson., J. Graham., M. M. Abou-Setta., El-Morshedy M. M, 1995. Estimating crop loss in orchards with patches of mature citrus trees infected by *Tylenchulus semipenetrans*. *Nematropica*. 25(1):43-51
- Duncan, L. W., (2005). Nematode parasites of Citrus. pp. 437–466. *In*: Plant Parasitic Nematodes in Subtropical and Tropical Agriculture (Luc M, Sikora RA, Bridge J, eds). *CAB International*, Wallingford. The UK.
- Fortuner R., A. R. Maggenti., L. M. Whittaker., 1984. Morphometrical variability in *Helicotylenchus* Steiner, 1945. 4: Study of field populations of *H. pseudorobustus* and related species. *Revue Nematologie*. 7: 121–135.
- Gaspard, J. T., R. Mankau., 1986. Nematophagous fungi associated with *Tylenchulus semipenetrans* and the citrus rhizosphere. *Nematologic*. 32: 359-363.
- Gottlieb Y., E. Cohn., Spiegel-Roy P, 1986. Biotypes of the citrus nematode (*Tylenchulus semipenetrans* Cobb) in Israel. *Phytoparasitica*. 14: 193-198.
- Gowen, S. R., P. Quénéhervé., R. Fogain., 2005. Nematode parasites of bananas and plantains. pp. 611-643. *In*: Plant Parasitic Nematodes in Subtropical and Tropical Agriculture. *CAB International*. Wallingford. The UK.
- Handoo, Z. A., A. M. Skantar., R. P. Mulrooney., 2010. First report of the sting nematode *Belonolaimus longicaudatus* on soybean in Delaware. *Plant Disease*. 94(1): 133.
- Hockland. S., R. N. Inserra., L. Millar., P. S. Lehman, 2006. International plant health- Putting legislation into practice. pp. 327–345. *In*: *Plant Nematology*. CAB International, Wallingford. The UK.
- Hollis, J. P. (1962). A survey of plant-parasitic nematodes and their control in Kenya. *FAO Plant Protection Bulletin*. 10: 97–106.
- Huettel, R. N., D. W. Dickson., 1981. Parthenogenesis in the two races of *Radopholus similis* from Florida. *Journal of Nematology*. 13: 13-15.
- Huettel, R. N., D. W. Dickson., D. T. Kaplan., 1983. Biochemical identification of the two races of *Radopholus similis* by starch gel electrophoresis. *Journal of Nematology*. 15: 338–344.

- Huettel, R. N., T. Yaegashi. (1988). Morphological differences between *Radopholus citrophilus* and *R. similis*. *Journal of Nematology*. 20: 150–157.
- Inserra, R. N., N. Vovlas. M Di Vito. (1994). Identification of second-stage juveniles of *Tylenchulus* spp. on the basis of posterior body morphology. *Nematropica*. 24: 25-33.
- Inserra, R. N., N. Vovlas., J. H. O'Bannon., R. P. Esser., 1988. *Tylenchulus graminis* n. sp. and *T. palustris* n. sp. (*Tylenchulidae*) from native flora of Florida, with notes on *T. semipenetrans* and *T. furcus*. *Journal of Nematology*. 20:266-287.
- Kaplan, D. T., (1986). Variation in *Radopholus citrophilus* population densities in the citrus rootstock Carrizo Citrange. *Journal of Nematology*. 18: 31–34.
- Kaplan, D. T., C. H. Opperman. (1997). Genome similarity implies that citrus-parasitic burrowing nematodes do not represent a unique species. *Journal of Nematology*. 29: 430–440.
- Kaplan, D. T., C. H. Opperman. 2000. Reproductive strategies and karyotype of the burrowing nematode, *Radopholus similis*. *Journal of Nematology*. 32: 126–133.
- Kaplan, D.T., J. H. O'Bannon. (1985). Occurrence of biotypes in *Radopholus citrophilus*. *Journal of Nematology*. 17: 158–162.
- Le Roux, H.F., (1995). Control of the citrus nematode in South Africa. PhD Dissertation, University of Pretoria, Pretoria, South Africa.45: 45–87.
- Loos, C. A., (1962). Studies on the life history and habits of the burrowing nematode, *Radopholus similis*, the cause of blackhead disease of banana. *Journal of Nematology*. 29: 43–52.
- McSorley R. (1986). Nematode problems on bananas and plantains in Florida. Nematology Circular No. 133, p 4. *Florida Department of Agriculture and Consumer Services*. 23: 34–67.
- McSorley R., J. L. Parrado., 1983. The spiral nematode *Helicotylenchus multicinctus* on bananas in Florida and its control. *Proceedings of the Florida State Horticultural Society*. 96: 201–207.
- McSorley R., J. L. Parrado. 1986. *Helicotylenchus multicinctus* on bananas: An international problem. *Nematropica*. 16: 73–91.
- Milne, D. L., P. Willers., 1979. Yield and nutritional responses to phenamiphos treatment of Citrus infested with citrus nematodes. *Subtropical*. 1:11-14.
- Mukhopadhyaya, M. C., D. Suryanarayana., 1969. Citrus decline in Haryana. The role of *Tylenchulus semipenetrans* and its control. *Indian Phytopathology*. 22:495-497.
- Nair MRGK., (1965). On the occurrence of the citrus nematode, *Tylenchulus semipenetrans* Cobb, in Kerala. *Indian Journal of Agricultural Research*. 3:105.
- O'Bannon, J. H., (1977). Worldwide dissemination of *Radophilus similis* and its importance in crop production. *Journal of Nematology*. 9: 16–25.
- O'Bannon, J. H., R. N. Inserra., 1989. *Helicotylenchus* species as crop-damaging parasitic nematodes. Nematology Circular 165. Florida Department of Agriculture and Consumer Services Division of Plant Industry.
- Pang W., J.E. Luc., W. T. Crow., K. Kenworthy., R. McSorly., J. K. Kruse., R. M. Giblin-Davis., 2011. Responses of seashore paspalum cultivars to sting and spiral nematodes. *Crop Science*. 51: 2864–2867.
- O'Bannon, J. H. (1968). The influence of an organic soil amendment on infectivity and reproduction of *Tylenchulus semipenetrans* on two citrus rootstocks. *Phytopathology*. 58: 597–601.
- Pang, W., J. E. Luc., W. T. Crow., K. E. Kenworthy., R. M. Giblin-Davis., J. K. McSorley, R, Kruse. 2012. Field responses of bermudagrass and seashore paspalum to sting and spiral nematodes. *Journal of Nematology*. 43: 201–208.
- Reddy, P. P., (1977). Plant-parasitic nematodes associated with Citrus in Karnataka, Kerala, Maharashtra, and Tamil Nadu states. (Abs). In: International Symposium of Citriculture, Horticultural Society of India, Bangalore, *Institute of Agricultural Research*. 39: 65–78.
- Reynolds, H. W., J. H. O'Bannon., 1963. Decline of grapefruit trees in relation to citrus nematode

- populations and tree recovery after chemical treatment. *Phytopathology*. 53: 1011–1015.
- Sher, S. A. (1968). Revision of the genus *Radopholus* Thorne, 1949 (Nematoda: *tylenchidia*). *Proceedings of the Helminthological Society of Washington*. 35, 219–237.
- Subbotin, S. A. R. N. Inserra., M. Marias., P. Mullin., T. O. Powers., P. A. Roberts., E. Van Den Berg., G.W. Yates., J.G. Baldwin., 2011. Diversity and phylogenetic relationships within the spiral nematodes of (*Helicotylenchus Steiner1945*) (*Tylenchida: Hoplolaimidae*) as inferred from analysis of the D2-D3 expansion segments of 28S rRNA gene sequences. *Nematology*. 13: 333–345
- Subbotin, S. A., N. Vovlas., G. W. Yeates., J. Hallmann., S. Kiewnick., V. N. Chizhov., Manzanilla-Lopez, R. H., R. N. Inserra., P. Castillo., 2015. Morphological and molecular characterization of *Helicotylenchus pseudorobustus*(Steiner, 2014) Golden, 1956 and related species (*Tylenchida: Holplolaimidae*) with a phylogeny of the genus. *Nematology*. 17: 27–52.
- Vu T. R. Hauschild R. A. Sikora. 2006. *Fusarium oxysporum* endophytes induced systemic resistance against *Radopholus similis* on bananas. *Nematology*. 8: 847-852.